

F1



Europäisches Patentamt
European Patent Office
Office européen des brevets



⑪ Publication number: **0 525 960 A1**

⑫

EUROPEAN PATENT APPLICATION

⑪ Application number: 92305287.2

⑤① Int. Cl. 5: A61K 35/74

⑫ Date of filing: 09.06.92

③① Priority: 18.06.91 US 717773

④③ Date of publication of application:
03.02.93 Bulletin 93/05

④④ Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL PT
SE

⑦① Applicant: AMERICAN HOME PRODUCTS
CORPORATION
685 Third Avenue
New York New York 10017-4085(US)

⑦② Inventor: Miller, Glenn Alvin

9780 NW 51st La.
Miami, Dade, Florida(US)
Inventor: Rabin, Mark Barry
8260 SW 146th Street
Miami, Dade, Florida(US)
Inventor: Harrington, William Joseph, Jnr.
9301 SW 92nd Avenue
Miami, Dade, Florida(US)

⑦④ Representative: Wileman, David Francis Dr. et
al
c/o Wyeth Laboratories Huntercombe Lane
South
Taplow Maidenhead Berkshire SL6 OPH(GB)

⑤④ Use of rapamycin for the treatment of adult T-cell leukemia/lymphoma.

⑤⑦ This invention provides a method of treating adult T-cell leukemia/lymphoma (ATL) in a mammal in need thereof which comprises administering an antiproliferative amount of rapamycin, and is particularly useful in arresting the progression of or inducing remission of ATL.

EP 0 525 960 A1

This invention relates to a method of treating adult T-cell leukemia/lymphoma, in particular to a new therapeutic use of rapamycin.

Adult T-cell Leukemia/Lymphoma (ATL) is an aggressive hematologic disease, characterized by a rapid, invariably fatal course. Clinical features of ATL include hypercalcemia, bone marrow infiltration, generalized lymphadenopathy, leukemia, skin involvement, hepatosplenomegaly and lytic bone lesions.

The human T-lymphotropic virus 1 (HTLV-1) is the etiologic agent of ATL. Evidence linking HTLV-1 to ATL includes epidemiologic clustering of ATL in geographic regions endemic for HTLV-1 infection [Hinuma, Y., Proc. Natl. Acad. Sci. 78: 6476 (1981); Watanabe, T., Virology 133: 238 (1984); Catovsky, D. Lancet 1: 639 (1982); Blattner, W.A., Int. J. Ca. 30: 257 (1982)], the high frequency of specific antibodies in sera from ATL patients, [Hinuma, Y., Proc. Natl. Acad. Sci. 78: 6476 (1981)], detection of monoclonally integrated proviral sequences in leukemic cells of patients with this disease [Yoshida M., Proc. Natl. Acad. Sci. 81: 2534 (1984)], and isolation of infectious virus from leukemia cell cultures [Poesz, B.J., Proc. Natl. Acad. Sci. 77: 7415 (1980); Yoshida, M., Proc. Natl. Acad. Sci. 79: 2031 (1982)].

HTLV-1 transformed T-cells are characterized by uncontrolled proliferation in the absence of antigenic stimulation. IL-2 independent growth is a hallmark of these cells in culture. The cell surface phenotype of these cells is dominated by a significant increase in IL-2 receptors. Recent evidence demonstrated that these receptors are high affinity IL-2R. The majority of T-cells isolated from infected individuals tested did not, however, respond to exogenously added IL-2. [Kodaka, T., Jpn. J. Ca. Res. 81: 902 (1990)].

HTLV-1 has been reported to be transmitted sexually, from mother-child via breast feeding, among intravenous drug abusers (presumably by sharing of contaminated needles), and via blood transfusion [Hino, S., Gann 76: 474 (1985); Murphy, E.L., Ann. Int. Med. 111: 555 (1989); Sato, H., Int. J. Ca. 37: 395 (1986); Robert-Guroff, M., J.A.M.A., 255: 3133 (1986)]. Infection with the virus has an associated lifetime risk for developing ATL estimated to be approximately 4% [Murphy, E.L., Ann. Int. Med. 111: 555 (1989)].

The prevalence of HTLV-1 infection in residents of the United States has recently become a public health concern. An epidemiologic survey undertaken in major cities of the United States detected average seroprevalence rates of 0.025 percent among healthy blood donors [Williams, A.E., Science, 240: 643 (1988)]. More recent epidemiologic data indicate a seropositivity rate of 4.4 percent among healthy blacks in Brooklyn, New York suggesting seroprevalence may vary significantly depending on the population sampled. [Dosik, H., Proc. Am. Soc. Clin. Oncol. 9: 2 (1990)]. Khabbaz [J.A.M.A. 263: 60 (1990)] reported average seroprevalence rates of 6.7% among prostitutes nationwide. These data ranged from a low of 0% in Nevada to a high of 25.4% in Newark, NJ indicating variable rates even among high risk groups.

Internationally, HTLV-1 is present in endemic proportions in several areas of the world. The southwestern islands of Japan have long been identified as an endemic region for HTLV-1 infection. [Jpn. J. Clin. Oncol. 15: 517 (1985)]. For example, seroprevalence rates as high as 8% have been detected in healthy volunteer blood donors in the southern province of Kyushu. [Maeda, Y., Int. J. Ca. 33: 717 (1984)]. A seroprevalence rate of 9.2% among otherwise healthy female food service workers over the age of 30 was reported in Jamaica. [Murphy, E.L., Ann. Int. Med. 111: 555 (1989)]. The seroprevalence rate rose to 14.1% in age matched females seen at a Jamaican sexually transmitted disease clinic. Risk factors for infection with the virus follow the expected pattern of a sexually transmitted disease.

Although serological and molecular genetic tests are available to screen for HTLV-1 infection, there is currently no means of early diagnosis of ATL. Presently, there is no effective therapeutic treatment for late stage disease, and as such mean survival after diagnosis is less than 6 months. Unsuccessful treatments were reported using radiation; methotrexate; leukovorin; interferon SQ; high dose steroids; and combinations containing cyclophosphamide, adriamycin, oncovin, prednisone, and bleomycin therapies. [Harrington, J. Aids 4: 284 (1991)]. Efficacious treatment of ATL was reported with SUN4800 (recombinant interferon gamma). [Ishihara, K., J. Invest. Dermatol. 93: 556 (1989)]. Responses of 90% were reported in 3 of 10 ATL patients treated with anti-Tac, a mouse antibody against the human p55 IL-2 receptor peptide (Tac; CD25). [Junghans, R., 37: 861A (1989)].

Rapamycin, a macrocyclic triene antibiotic produced by *Streptomyces hygroscopicus* [U.S. Patent 3,929,992] has been shown to prevent the formation of humoral (IgE-like) antibodies in response to an albumin allergic challenge [Martel, R., Can. J. Physiol. Pharm. 55: 48 (1977)], inhibit murine T-cell activation [Strauch, M., FASEB 3: 3411 (1989)], and prolong survival time of organ grafts in histoincompatible rodents [Morris, R., Med. Sci. Res. 17: 877 (1989)].

This invention provides a method of treating adult T-cell leukemia/lymphoma (ATL) in a mammal in need thereof which comprises administering an antiproliferative amount of rapamycin orally, parenterally, intranasally, intrabronchially, topically, transdermally, or rectally. In particular, rapamycin is useful in arresting the progression of and inducing remission of ATL by virtue of its ability to inhibit the proliferation of HTLV-1 transformed T-cells.

This invention also provides use of rapamycin in the preparation of a medicament for treating adult T-cell leukemia/lymphoma in a mammal.

The effect of rapamycin on ATL was established in an *in vitro* standard pharmacological test procedure using ten human HTLV-1 transformed T-cell lines cultured from ATL patients. These cell lines are presently the best models for ATL in humans. Cyclosporin A (CsA) was also evaluated for the purpose of comparison. The procedure used and results obtained are described below.

ATL cell lines were established according to known literature procedures. ATL5S, ATL3I, and ATL1K were established according to the method of Hoshino [Proc. Natl. Acad. Sci, 80: 6061 (1983)]. The cell lines 31008, 31009, 31016 and AT(EGG) were established from peripheral blood of HTLV-1 infected individuals via co-culture with cord blood lymphocytes. The cell line MALN was isolated from lymphoid tumor tissue of an HTLV-1 infected ATL patient seen at Jackson Memorial/University of Miami Medical Center. [Harrington, W.J., Jr., J. Acq. Imm. Def. Dis., 4: 284 (1991)]. The primary cultures BL and GP were initiated from peripheral blood of ATL patients. The MALN, BL, and GP cell lines were initiated without cultivation.

Cells were adjusted to a concentration of 1×10^6 cells/mL in RPMI 1640 with 15% fetal bovine serum and were supplemented with 50 U/mL penicillin, 50 μ g/mL streptomycin, and 2 mM glutamine. Rapamycin and CsA were each diluted to a stock concentration of 1 mM in absolute ethanol with subsequent dilutions being made in culture medium. Drug concentrations evaluated ranged from 1 μ M to 100 fM. Cultures were incubated at 37°C, 5% CO₂ for 24 hours followed by a 4 hr. pulse with 0.5 μ Ci of tritiated thymidine. Thymidine incorporation was measured by liquid scintillation counting, and the results expressed in counts per minute (cpm). The results obtained from the mean of four evaluations at each concentration were compared with control values to determine percent inhibition. The following formula was used to calculate the percent inhibition:

$$\text{Percent Inhibition} = \frac{\text{cpm}[\text{no drug}] - \text{cpm}[\text{drug}]}{\text{cpm}[\text{no drug}]} \times 100$$

The percent inhibition obtained with rapamycin is shown in the following table.

PERCENT INHIBITION OF ATL CELL PROLIFERATION										
Rapamycin Concentration	Cell Line									
	ATL3I	ATL5S	ATL1K	ATEGG	31008	31009	31016	MALN	GP	BL
1 μ M	28	59	19	58	69	71	59	55	54	80
100 nM	17	58	32	60	68	70	56	54	46	79
10 nM	15	49	18	45	58	67	49	41	46	76
1 nM	18	48	26	53	54	68	48	43	42	73
100 pM	11	47	23	49	45	61	49	37	39	65
10 pM	6	38	12	44	31	47	47	26	14	55
1 pM	11	14	15	48	30	59	50	36	20	56
100 fM	7	36	15	44	29	60	44	36	23	51

Rapamycin inhibited proliferation of 8 of the 10 ATL cell lines at a concentration of 100 nM or greater, with a 50% reduction in proliferation in 4 of the cell lines at a concentration of 1 nM. Both of the cell lines in which proliferation was not inhibited by rapamycin had been in culture for more than 5 years without recloning; their nonresponsiveness to rapamycin may be attributed to their further transformation into insensitive clones during this time period. In comparison, CsA did not inhibit proliferation in 7 of the 10 cell lines in which it was evaluated. In the remaining 3 cell lines (ATL3I, ATL5S, and 31009) proliferation was inhibited by 16%, 23%, and 22%, respectively.

The results obtained in this standard pharmacological test procedure demonstrate that rapamycin inhibits the proliferation of HTLV-1 transformed T-cell lines, which are presently the best models for ATL in humans, and as such, rapamycin is useful in the treatment of ATL.

When rapamycin is employed in the treatment of ATL, it can be formulated neat or with a pharmaceutical carrier to a mammal in need thereof preferably in unit dosage form; conveniently a single unit dose for daily administration. The pharmaceutical carrier may be solid or liquid.

A solid carrier can include one or more substances which may also act as flavoring agents, lubricants,

solubilizers, suspending agents, fillers, glidants, compression aids, binders or tablet-disintegrating agents; it can also be an encapsulating material. In powders, the carrier is a finely divided solid which is in admixture with the finely divided active ingredient. In tablets, the active ingredient is mixed with a carrier having the necessary compression properties in suitable proportions and compacted in the shape and size desired. Suitable solid carriers include, for example, calcium phosphate, magnesium stearate, talc, sugars, lactose, dextrin, starch, gelatin, cellulose, methyl cellulose, sodium carboxymethyl cellulose, polyvinylpyrrolidone, low melting waxes and ion exchange resins.

Liquid carriers are used in preparing solutions, suspensions, emulsions, syrups, elixirs and pressurized compositions. The active ingredient can be dissolved or suspended in a pharmaceutically acceptable liquid carrier such as water, an organic solvent, a mixture of both or pharmaceutically acceptable oils or fats. The liquid carrier can contain other suitable pharmaceutical additives such as solubilizers, emulsifiers, buffers, preservatives, sweeteners, flavoring agents, suspending agents, thickening agents, colors, viscosity regulators, stabilizers or osmo-regulators. Suitable examples of liquid carriers for oral and parenteral administration include water (partially containing additives as above, e.g. cellulose derivatives, preferably sodium carboxymethyl cellulose solution), alcohols (including monohydric alcohols and polyhydric alcohols, e.g. glycols) and their derivatives, and oils (e.g. fractionated coconut oil and arachis oil). For parenteral administration, the carrier can also be an oily ester such as ethyl oleate and isopropyl myristate. Sterile liquid carriers are useful in sterile liquid form compositions for parenteral administration. The liquid carrier for pressurized compositions can be halogenated hydrocarbon or other pharmaceutically acceptable propellant.

Liquid pharmaceutical compositions which are sterile solutions or suspensions can be utilized by, for example, intramuscular, intraperitoneal or subcutaneous injection. Sterile solutions can also be administered intravenously. The compound can also be administered orally either in liquid or solid composition form.

Rapamycin may be administered rectally in the form of a conventional suppository. For administration by intranasal or intrabronchial inhalation or insufflation, rapamycin may be formulated into an aqueous or partially aqueous solution, which can then be utilized in the form of an aerosol. Rapamycin may also be administered transdermally through the use of a transdermal patch containing the active compound and a carrier that is inert to the active compound, is non-toxic to the skin, and allows delivery of the agent for systemic absorption into the blood stream via the skin. The carrier may take any number of forms such as creams and ointments, pastes, gels, and occlusive devices. The creams and ointments may be viscous liquid or semisolid emulsions of either the oil-in-water or water-in-oil type. Pastes comprised of absorptive powders dispersed in petroleum or hydrophilic petroleum containing the active ingredient may also be suitable. A variety of occlusive devices may be used to release the active ingredient into the blood stream such as a semipermeable membrane covering a reservoir containing the active ingredient with or without a carrier, or a matrix containing the active ingredient. Other occlusive devices are known in the literature.

Rapamycin may be administered topically as a solution, cream, or lotion by formulation with pharmaceutically acceptable vehicles containing 0.1 - 5 percent, preferably 2%, of active compound.

The dosage requirements vary with the particular compositions employed, the route of administration, the severity of the symptoms presented and the particular subject being treated. Based on the results obtained in the standard pharmacological test procedure, projected parenteral daily dosages of active compound would be 0.01 $\mu\text{g/kg}$ - 10 mg/kg and preferably between 0.1 $\mu\text{g/kg}$ - 2 mg/kg. When rapamycin is given parenterally, it will preferably be administered intravenously. Projected oral daily dosages of active compound would be 1 $\mu\text{g/kg}$ - 15 mg/kg and preferably between 0.01 mg/kg - 10 mg/kg. Treatment will generally be initiated with small dosages less than the optimum dose of the compound. Thereafter the dosage is increased until the optimum effect under the circumstances is reached; precise dosages for oral, parenteral, nasal, or intrabronchial administration will be determined by the administering physician based on experience with the individual subject treated. In general, rapamycin is most desirably administered at a concentration that will generally afford effective results without causing any harmful or deleterious side effects, and can be administered either as a single unit dose, or if desired, the dosage may be divided into convenient subunits administered at suitable times throughout the day.

For the treatment of ATL, it is also contemplated that rapamycin may be administered in conjunction with chemotherapeutic agents such as methotrexate, leukovorin, cyclophosphamide, adriamycin, oncovin, prednisone, bleomycin, and the like; high dose steroids; radiation therapy; cytokine therapy, such as interferon; monoclonal antibody therapy; and other therapies useful in the treatment of neoplastic diseases.

Claims

1. Use of rapamycin in the preparation of a medicament for treating adult T-cell leukemia/lymphoma in a

mammal.

2. Use of rapamycin according to Claim 1 in which the medicament is adapted for administration orally, parenterally, intranasally, intrabronchially, topically, transdermally or rectally.
- 5 3. Use of rapamycin according to Claim 2 in which the medicament is adapted for intravenous administration.
- 10 4. Use of rapamycin according to Claim 3 in which the medicament is in unit dosage form to provide a daily dosage from 0.01 $\mu\text{g/kg}$ to 10 mg/kg based on the weight of the mammal to treated.
5. Use of rapamycin according to Claim 3 in which the medicament is in unit dosage form to provide a daily dosage from 0.1 $\mu\text{g/kg}$ to 2 mg/kg based on the weight of the mammal to treated.
- 15 6. Use of rapamycin according to Claim 1 in which the medicament is adapted for oral administration.
7. Use of rapamycin according to Claim 6 in which the medicament is in unit dosage form to provide a daily dosage from 1 $\mu\text{g/kg}$ to 15 mg/kg based on the weight of the mammal to treated.
- 20 8. Use of rapamycin according to Claim 6 in which the medicament is in unit dosage form to provide a daily dosage from 0.01 mg/kg to 10 mg/kg based on the weight of the mammal to treated.

25

30

35

40

45

50

55



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 92 30 5287

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
P, X	THE JOURNAL OF IMMUNOLOGY, vol. 148, no. 10, 15th May 1992, pages 3256-3263, The American Association of Immunologists, US; P. HÖLLSBERG et al.: "Characterization of HTLV-I in vivo infected T cell clones. IL-2-independent growth of nontransformed T cells" * Whole document *	1	A 61 K 35/74
A	THE JOURNAL OF IMMUNOLOGY, vol. 144, no. 4, 15th February 1990, pages 1418-1424, The American Association of Immunologists, US; F.J. DUMONT et al.: "The immunosuppressive macrolides FK-506 and rapamycin act as reciprocal antagonists in murine T cells" * Abstract; page 1423, lines 42-53 *	1-8	
A	TRANSPLANTATION PROCEEDINGS, vol. 23, no. 2 (suppl. 2), 1991, pages 1-5, PROCEEDINGS OF A SATELLITE SYMPOSIUM ON IMMUNOSUPPRESSIVE DRUGS: TARGETS AND MECHANISMS OF ACTION, San Francisco, CA, 19th August 1990; N.H. SIGAL et al.: "Inhibition of human T-cell activation by FK 506, rapamycin, and cyclosporine A" * Whole document *	1-8	TECHNICAL FIELDS SEARCHED (Int. Cl.5) A 61 K
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 28-09-1992	Examiner MAIR J.
CATEGORY OF CITED DOCUMENTS X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document		T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons A: member of the same patent family, corresponding document	



European Patent
Office

EUROPEAN SEARCH REPORT

Page 2

Application Number

EP 92 30 5287

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
P,X	JOURNAL OF NEUROIMMUNOLOGY, vol. 0, suppl. 1, 1991, page 201, 3RD INTERNATIONAL CONFERENCE ON NEUROIMMUNOLOGY, Jerusalem, 27th October - 1st November 1991; P. HÖLLSBERG et al.: "HTLV-1 induced spontaneous T cell clonal proliferation is rapamycin sensitive" * Whole document *	1	
A	TRENDS IN PHARMACOLOGICAL SCIENCES, vol. 12, no. 6, June 1991; pages 218-223, Elsevier Science Publishers Ltd, GB; J.Y. CHANG et al.: "FK-506 and rapamycin: novel pharmacological probes of the immune response" * Whole document *	1-8	
A	BE-A- 877 700 (AYERST, McKENNA & HARRISON, INC.) * Whole document *	1-8	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
Place of search THE HAGUE		Date of completion of the search 28-09-1992	Examiner MAIR J.
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons * : member of the same patent family, corresponding document			

EP FORM 150 (3.91) (P0001)